

**Literaturverzeichnis zum Fachartikel „Die größten Irrtümer in der Neuroreha oder wie Neurorehabilitation effektiv wird“ von Sabine und Hans Lamprecht in „praxis ergotherapie“, Ausgabe 03-2024:**

- Ada, L.; Dorsch, S.; Canning, C. G. (2006): Strengthening interventions increase strength and improve activity after stroke: a systematic review. In: *The Australian journal of physiotherapy* 52 (4), S. 241–248 .
- Ada, L.; Vattanasilp, W.; O'Dwyer, N. J.; Crosbie, J. (1998): Does spasticity contribute to walking dysfunction after stroke? In: *Journal of neurology, neurosurgery, and psychiatry* 64 (5).
- Ardestani, M M.; Kinnaird, C R.; Henderson, Ch E.; Hornby, T. G (2019): Compensation or Recovery? Altered Kinetics and Neuromuscular Synergies Following High-Intensity Stepping Training Poststroke. In: *Neurorehabilitation and neural repair* 33 (1).
- Blum, Carlos González; Scherfer, Erwin (2023): Neuro-Reha. In: *physiopraxis* 21 (02).
- Canning, C G.; Ada, L; Adams, R; O'Dwyer, N J. (2004): Loss of strength contributes more to physical disability after stroke than loss of dexterity. In: *Clinical rehabilitation* 18.
- Carr J H; Shepherd R B: *Stroke Rehabilitation* (2003): Elsevier.
- Chiviacowsky, Suzete; Wulf, Gabriele; Wally, Raquel; Borges, Thiago (2009): Knowledge of results after good trials enhances learning in older adults. In: *Research quarterly for exercise and sport* 80 (3), S. 663–668.
- Crossman, E. R. F. W. (1959): A theory of the acquisition of speed-skill. In: *Ergonomics* 2 (2), S. 153–166.
- Dickstein, R.; Heffes, Y.; Laufer, Y.; Ben-Haim, Z. (1999): Activation of selected trunk muscles during symmetric functional activities in poststroke hemiparetic and hemiplegic patients. In: *Journal of neurology, neurosurgery, and psychiatry* 66 (2), S. 218–221.
- Dobkin, B H. (2005): Clinical practice. Rehabilitation after stroke. In: *The New England journal of medicine* 352 (16).
- Hillier, S; Worley, A. (2023): The effectiveness of the feldenkrais method. A systematic review of the evidence. *Feldenkrais Research Journal*. 7, (Apr. 2023).
- Högg, Susan (2023): Armkraftraining in der Schlaganfallrehabilitation. In: *neuroreha* 15 (02), S. 66–72.
- Huber, Martin (2022): Das „Wie“ ist entscheidend – Die Renaissance der Bewegungsqualität. In: *physiopraxis* 20 (09).

Kim, Nan-Hyang; Cha, Yong-Jun (2015): Effect of gait training with constrained-induced movement therapy (CIMT) on the balance of stroke patients. In: *Journal of physical therapy science* 27 (3), S. 611–613.

Kirker, S. G.; Simpson, D. S.; Jenner, J. R.; Wing, A. M. (2000): Stepping before standing: hip muscle function in stepping and standing balance after stroke. In: *Journal of neurology, neurosurgery, and psychiatry* 68 (4), S. 458–464.

Lamprecht S. und Lamprecht H. CIMT - Constraint Induced Movement Therapy - Evidenzbasierte Therapie bei leichter und mittlerer Armparese. Springer Nature (in Druck)

Lamprecht S.; Lamprecht H.: „Die zehn größten Irrtümer in der Neuroreha“; PT Zeitschrift für Physiotherapeuten ; 09/2021; Pflaumverlag

Mahtani, G B.; Kinnaird, C R.; Connolly, M; Holleran, C L.; Hennessy, P W.; Woodward, J et al. (2017): Altered Sagittal- and Frontal-Plane Kinematics Following High-Intensity Stepping Training Versus Conventional Interventions in Subacute Stroke. In: *Physical therapy* 97 (3).

Majsak, Michael J. (1996): Application of Motor Learning Principles to the Stroke Population. In: *Topics in stroke rehabilitation* 3 (2), S. 37–59.

Nikamp, C; Buurke, J; Schaake, L; van der Palen, J; Rietman, J; Hermens, H (2019): Effect of long-term use of ankle-foot orthoses on tibialis anterior muscle electromyography in patients with sub-acute stroke: A randomized controlled trial. In: *Journal of rehabilitation medicine* 51 (1).

Oujamaa, L.; Relave, I.; Froger, J.; Mottet, D.; Pelissier, J-Y (2009): Rehabilitation of arm function after stroke. Literature review. In: *Annals of physical and rehabilitation medicine* 52 (3).

Pak, Sang; Patten, Carolynn (2008): Strengthening to promote functional recovery poststroke: an evidence-based review. In: *Topics in stroke rehabilitation* 15 (3).

Park, Seoung Hoon; Hsu, Chao-Jung; Dee, Weena; Roth, Elliot J.; Rymer, William Z.; Wu, Ming (2021): Enhanced error facilitates motor learning in weight shift and increases use of the paretic leg during walking at chronic stage after stroke. In: *Experimental brain research* 239 (11), S. 3327–3341.

Platz T. et al., Therapie des spastischen Syndroms, S2k-Leitlinie, 2018, in: Deutsche Gesellschaft für Neurologie (Hrsg.), Leitlinien für Diagnostik und Therapie in der Neurologie. Online: [www.dgn.org/leitlinien](http://www.dgn.org/leitlinien) (abgerufen am 14.08.2023)

S3-Leitlinie „Rehabilitative Therapie bei Armparese nach Schlaganfall“ der DGKR

Souza-Teixeira, F. de; Costilla, S.; Ayán, C.; García-López, D.; González-Gallego, J.; Paz, J. A. de (2009): Effects of Resistance Training in Multiple Sclerosis. In: *Int J Sports Med* 30 (04).

Sterr, A; Freivogel, S. (2004): Intensive training in chronic upper limb hemiparesis does not increase spasticity or synergies. In: *Neurology* 63 (11), S. 2176–2177.

Taub, E. (1976): Movement in nonhuman primates deprived of somatosensory feedback. In: *Exercise and sport sciences reviews* 4, S. 335–374.

Thijs, L; Denissen, S; Mehrholz, J; Elsner, B; Lemmens, R; Verheyden, G; (2020): Trunk training for improving activities in people with stroke. In: *Cochrane Database of Systematic Reviews*. DOI: 10.1002/14651858.CD013712.

Wagner, J M.; Lang, C E.; Sahrman, S A.; Edwards, D F.; Dromerick, A W. (2007): Sensorimotor impairments and reaching performance in subjects with poststroke hemiparesis during the first few months of recovery. In: *Physical therapy* 87 (6), S. 751–765.

Wens, I.; Dalgas, U.; Vandenabeele, F.; Grevendonk, L.; Verboven, K.; Hansen, D.; Eijnde, B O. (2015): High Intensity Exercise in Multiple Sclerosis: Effects on Muscle Contractile Characteristics and Exercise Capacity, a Randomised Controlled Trial. In: *PloS one* 10 (9), e0133697.

White, L. J.; McCoy, S. C.; Castellano, V.; Gutierrez, G.; Stevens, J. E.; Walter, G. A.; Vandenborne, K. (2004): Resistance training improves strength and functional capacity in persons with multiple sclerosis. In: *Mult Scler* 10 (6).

Wong, Y; Ada, L; Månum, G; Langhammer, B (2023): Upper limb practice with a dynamic hand orthosis to improve arm and hand function in people after stroke: a feasibility study. In: *Pilot and feasibility studies* 9 (1),